

This text is meant purely as a documentation tool and has no legal effect. The Union's institutions do not assume any liability for its contents. The authentic versions of the relevant acts, including their preambles, are those published in the Official Journal of the European Union and available in EUR-Lex. Those official texts are directly accessible through the links embedded in this document

► **B**

COMMISSION REGULATION (EC) No 1847/2003

of 20 October 2003

concerning the provisional authorisation of a new use of an additive and the permanent authorisation of an additive already authorised in feedingstuffs

(Text with EEA relevance)

(OJ L 269, 21.10.2003, p. 3)

Amended by:

		Official Journal		
		No	page	date
► <u>M1</u>	Commission Implementing Regulation (EU) No 1018/2012 of 5 November 2012	L 307	56	7.11.2012
► <u>M2</u>	Commission Implementing Regulation (EU) 2018/347 of 5 March 2018	L 67	21	9.3.2018

▼B

COMMISSION REGULATION (EC) No 1847/2003

of 20 October 2003

**concerning the provisional authorisation of a new use of an additive
and the permanent authorisation of an additive already authorised
in feedingstuffs**

(Text with EEA relevance)

Article 1

The additive belonging to the group ‘enzymes’ referred to in Annex I is authorised for use as an additive in feedingstuffs under the conditions laid down in that Annex.

▼M2

▼B

Article 3

This Regulation shall enter into force on the third day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

▼ B

ANNEX I

No (or EC No)	Additive	Chemical formula, description	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
					Units of activity/kg of complete feedingstuff			
Enzymes								
14	Endo-1,4-beta-xylanase EC 3.2.1.8.	Preparation of endo-1,4-beta-xylanase produced by <i>Aspergillus niger</i> (CBS 520.94) having a minimum activity of: Solid form: 600 U/g ⁽¹⁾ Liquid form: 300 U/ml	Turkeys for fattening	—	300 U	—	1. In the directions for use of the additive and premixture, indicate the storage temperature, storage life, and stability to pelleting 2. Recommended dose per kg of complete feedingstuff: 300 — 1 200 U 3. For use in compound feed rich in non-starch polysaccharides, (mainly arabinoxylans) e.g. containing more than 40 % wheat.	24.10.2007

⁽¹⁾ One U is the amount of enzyme which liberates 1 micromole of xylose from birchwood xylan per minute at pH 5,3 and 50 °C.’

▼ M2