Commission Regulation (EC) No 1441/2007 of 5 December 2007 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs (Text with EEA relevance)

COMMISSION REGULATION (EC) No 1441/2007

of 5 December 2007

amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs

(Text with EEA relevance)

THE COMMISSION OF THE EUROPEAN COMMUNITIES,

Having regard to the Treaty establishing the European Community,

Having regard to Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004 on the hygiene of foodstuffs⁽¹⁾, and in particular Article 4(4) thereof,

Whereas:

- (1) Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs⁽²⁾ lays down microbiological criteria for certain micro-organisms and the implementing rules to be complied with by food business operators when implementing the general and specific hygiene measures referred to in Article 4 of Regulation (EC) No 852/2004. Regulation (EC) No 2073/2005 also provides that food business operators are to ensure that foodstuffs comply with the relevant microbiological criteria set out in Annex I to that Regulation.
- (2) Chapters 1 and 2 of Annex I to Regulation (EC) No 2073/2005 set out food safety criteria and process hygiene criteria regarding dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age (dried infant formulae and dried dietary foods). Part 2.2 of Chapter 2 of that Annex provides that where dried infant formulae and dried dietary foods are tested and Enterobacteriaceae are detected in any of the sample units, the batch is to be tested for *Enterobacter sakazakii* and *Salmonella*.
- (3) On 24 January 2007, the Scientific Panel on Biological Hazards (BIOHAZ Panel) of the European Food Safety Authority (EFSA) issued an opinion with regard to Enterobacteriaceae as indicators of *Salmonella* and *Enterobacter sakazakii*. It concluded that it is not possible to establish a correlation between Enterobacteriaceae and *Salmonella*, and no universal correlation between Enterobacteriaceae and *Enterobacter sakazakii* exists. At individual plant level, a correlation between Enterobacteriaceae and *Enterobacter sakazakii* exists.
- (4) Therefore the requirement laid down in Regulation (EC) No 2073/2005 as regards the testing of dried infant formulae and dried dietary foods for *Salmonella* and *Enterobacter* sakazakii where Enterobacteriaceae are detected in any of the sample units should no

longer apply. Part 2.2 of Chapter 2 of Annex I to that Regulation should therefore be amended accordingly.

- (5) In line with the opinion on the microbiological risks in infant formulae and follow-on formulae issued by the BIOHAZ Panel of EFSA on 9 September 2004, microbiological criteria on *Salmonella* and Enterobacteriaceae should be laid down for dried follow-on formulae.
- (6) The BIOHAZ Panel of EFSA issued an opinion on *Bacillus cereus* and other *Bacillus* spp. in foodstuffs on 26 and 27 January 2005. It concluded that one of the major control measures is to control temperature and to establish a system based on hazard analysis and critical control point principles. Dehydrated foods, in which the presence of spores of pathogenic *Bacillus* spp. is frequent, might permit the growth of *Bacillus cereus* once rehydrated in warm water. Some dehydrated foods, including dried infant formulae and dried dietary foods, are consumed by potentially fragile consumers. In line with the EFSA opinion, the numbers of *Bacillus cereus* spores in dried infant formulae and dried dietary foods should be as low as possible during processing and a process hygiene criterion should be laid down in addition to good practices designed to reduce delay between preparation and consumption.
- (7) Chapter 1 of Annex I to Regulation (EC) No 2073/2005 provides for the analytical reference method for staphylococcal enterotoxins in certain cheeses, milk powder and whey powder. That method has been revised by the Community reference laboratory for coagulase positive staphylococci. The reference to that analytical reference method should therefore be amended. Chapter 1 of Annex I to that Regulation should therefore be amended accordingly.
- (8) Chapter 3 of Annex I to Regulation (EC) No 2073/2005 sets out sampling rules for carcasses of cattle, pig, sheep, goats and horses for *Salmonella* analyses. Pursuant to those rules the sampling area is to cover a minimum of 100 cm² per site selected. However, neither the number of sampling sites nor the minimum total area of sampling is specified. In order to improve the implementation of these rules in the Community, it is appropriate to further specify in Regulation (EC) No 2073/2005 that the areas most likely to be contaminated should be selected for sampling and that the total sampling area should be increased. Chapter 3 of Annex I to that Regulation should therefore be amended accordingly.
- (9) In the interests of clarity of Community legislation, it is appropriate to replace Annex I to Regulation (EC) No 2073/2005 by the text set out in the Annex to this Regulation.
- (10) The measures provided for in this Regulation are in accordance with the opinion of the Standing Committee on the Food Chain and Animal Health,

HAS ADOPTED THIS REGULATION:

Article 1

Annex I to Regulation (EC) No 2073/2005 is replaced by the text in the Annex to this Regulation.

Article 2

This Regulation shall enter into force on the 20th day following its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 5 December 2007.

For the Commission Markos KYPRIANOU Member of the Commission

ANNEX

ANNEX I

Microbiological criteria for foodstuffs

Chapter 1.

Food safety criteria

Food	Micro-	Sampling plan ^a		Limits ^b		Analytical Stage	
category	organism their toxins, metabolit		c	m	М	reference method ^c	where the criterion applies
	<i>Listeria</i> Ready- to- eat foods intended for infants and ready- to- eat foods for special medical purposes ^d	10 enes	0	Absence in	25 g	EN/ISO 11290-1	Products placed on the market during their shelf-life
	<i>Listeria</i> Read <i>biocytog</i> to- eat foods able to support	5 enes	0	100 cfu/g ^e		EN/ISO 11290-2 ^f	Products placed on the market during their shelf-life
	the growth of <i>L. monocytogen</i> other than those intended for	5 es,	0	Absence in	25 g ^g	EN/ISO 11290-1	Before the food has left the immediate control of the food business operator, who has

Status: Point in time view as at 05/12/2007.

	infants and for special medical purposes					produced it
1.3	<i>Listeria</i> Read/ <i>hocytog</i> to- eat foods unable to support the growth of <i>L.</i> <i>monocytogen</i> other than those intended for infants and for special medical purposes ^{dh}		0	100 cfu/g	EN/ISO 11290-2 ^r	Products placed on the market during their shelf-life
1.4	Salmonella Minced meat and meat preparations intended to be eaten raw	25	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.5	Salmonella Minced meat and meat preparations made from poultry meat intended	15	0	From 1.1.2006 Absence in 10 g From 1.1.2010 Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life

	to					
	be					
	eaten					
	cooked					
1.6	Salmonella Minced meat	5	0	Absence in 10 g	EN/ISO 6579	Products placed on the
	and					market
	meat					during
	preparations					their
	made					shelf-life
	from					
	other					
	species					
	than					
	poultry intended					
	to					
	be					
	eaten					
	cooked					
		<i></i>	0	A1 10		
1.7	Salmonella Mechanically	5	0	Absence in 10 g	EN/ISO 6579	Products
	separated				03/9	placed on the
	meat					market
	(MSM) ⁱ					during
						their
						shelf-life
	Salmonalla	5	0	Absence in 25 g	EN/ISO	Products
1.8	<i>Salmonella</i> Meat	5	0	Absence in 25 g	6579	placed
	products				0377	on the
	intended					market
	to					during
	be					their
	eaten					shelf-life
	raw,					
	excluding products					
	where					
	the					
	manufacturin	g				
	process					
	or					
	the					
	composition					
	of					
	the					
	product					
	will eliminate					
	the					
	uic					

1.9	salmonella risk Salmonella Meat products made from poultrymeat intended to be eaten	5	0	From 1.1.2006 Absence in 10 g From 1.1.2010 Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.10	cooked Salmonella Gelatine and collagen	15	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.11	Salmonella Cheeses, butter and cream made from raw milk or milk that has undergone a lower heat treatment than pasteurisation		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.12	Salmonella Milk powder and whey powder	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.13	Salmonella Ice cream ^k , excluding	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market

	products where the manufacturin process or the composition of the product will eliminate the salmonella risk	g				during their shelf-life
1.14	Salmonella Egg products, excluding products where the manufacturing process or the composition of the product will eliminate the salmonella risk		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.15	Salmonella Ready- to- eat foods containing raw egg, excluding products where the manufacturin process or the		0	Absence in 25 g or ml	EN/ISO 6579	Products placed on the market during their shelf-life

	composition of the product will eliminate the salmonella risk					
1.16	Salmonella Cooked crustaceans and molluscan shellfish	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.17	Salmonella Live bivalve molluscs and live echinoderms, tunicates and gastropods		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.18	Salmonella Sprouted seeds (ready- to- eat) ¹	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.19	Salmonella Precut fruit and vegetables (ready- to- eat)	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.20	Salmonella Unpasteurised fruit and vegetable juices (ready- to- eat)	5 1	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life

Status: Point in time view as at 05/12/2007.

Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 1441/2007. (See end of Document for details)

1.21	Staphyloco Chemesestorio milk powder and whey powder, as referred to in the coagulase- positive staphylococci criteria in Chapter 2.2 of this Annex	S	0	Not detected in 25 g	European screening method of the CRL for coagulase positive staphylocod	Products placed on the market during their shelf-life cci ^m
1.22	Salmonella Dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age	30	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.23	<i>Salmonella</i> Dried follow- on formulae	30	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.24	<i>Enterobact</i> Dried infant	B 0	0	Absence in 10 g	ISO/TS 22964	Products placed on the

Status: Point in time view as at 05/12/2007.

			: There are curre n (EC) No 1441/				
	formulae and dried dietary foods for special medical purposes intended for infants below six months of						market during their shelf-life
1.25	age ⁿ <i>E. coli</i> ^o bivalve molluscs and live echinoderms, tunicates and gastropods	1 ^p	0	230 MPN/100 g of flesh and intra- valvular liquid		ISO TS 16649-3	Products placed on the market during their shelf-life
1.26	Histamine Fishery products from fish species associated with a high amount of histidine ^q	9 ^r	2	100 mg/ kg	200 mg/ kg	HPLC ^s	Products placed on the market during their shelf-life
1.27	Histamine Fishery products which have undergone enzyme maturation treatment in brine, manufactured		2	200 mg/ kg	400 mg/ kg	HPLC ^s	Products placed on the market during their shelf-life

	from fish species associated with a high amount of histidine ^q								
a	n = number of units comprising the sample; $c =$ number of sample units giving values between m and M.								
b	For points 1.1-1.25 m = M.								
c	The most recent edition of the standard shall be used.								
d	 Regular testing against the criterion is not required in normal circumstances for the following ready-to-eat foods: those which have received heat treatment or other processing effective to eliminate <i>L. monocytogenes</i>, when recontamination is not possible after this treatment (for example, products heat treated in their final package), fresh, uncut and unprocessed vegetables and fruits, excluding sprouted seeds, bread, biscuits and similar products, bottled or packed waters, soft drinks, beer, cider, wine, spirits and similar products, sugar, honey and confectionery, including cocoa and chocolate products, live bivalve molluscs. 								
e	This criterion shall apply if the manufacturer is able to demonstrate, to the satisfaction of the competent authority, that the product will not exceed the limit 100 cfu/g throughout the shelf-life. The operator may fix intermediate limits during the process that must be low enough to guarantee that the limit of 100 cfu/g is not exceeded at the end of shelf-life.								
f	1 ml of inoculum is plated on a Petri dish of 140 mm diameter or on three Petri dishes of 90 mm diameter.								
g	This criterion shall apply to products before they have left the immediate control of the producing food business operator, when he is not able to demonstrate, to the satisfaction of the competent authority, that the product will not exceed the limit of 100 cfu/g throughout the shelf-life.								
h	Products with $pH \le 4,4$ or $a_w \le 0,92$, products with $pH \le 5,0$ and $a_w \le 0,94$, products with a shelf-life of less than five days shall be automatically considered to belong to this category. Other categories of products can also belong to this category, subject to scientific justification.								
i	This criterion shall apply to mechanically separated meat (MSM) produced with the techniques referred to in paragraph 3 of Chapter III of Section V of Annex III to Regulation (EC) No 853/2004 of the European Parliament and of the Council.								
j	Excluding products when the manufacturer can demonstrate to the satisfaction of the competent authorities that, due to the ripening time and a_w of the product where appropriate, there is no salmonella risk.								
k	Only ice creams containing milk ingredients.								
1	Preliminary testing of the batch of seeds before starting the sprouting process or the sampling must be carried out at the stage where the highest probability of finding Salmonella is expected.								
m	<i>Reference:</i> Community reference laboratory for coagulase positive staphylococci. European screening method for the detection of staphylococcal enterotoxins in milk and milk products.								
n	Parallel testing for Enterobacteriaceae and <i>E. sakazakii</i> shall be conducted, unless a correlation between these micro- organisms has been established at an individual plant level. If Enterobacteriaceae are detected in any of the product samples tested in such a plant, the batch must be tested for <i>E. sakazakii</i> . It shall be the responsibility of the manufacturer to demonstrate to the satisfaction of the competent authority whether such a correlation exists between Enterobacteriaceae and <i>E. sakazakii</i> .								
0	<i>E. coli</i> is used here as an indicator of faecal contamination.								
р	A pooled sample comprising a minimum of 10 individual animals.								
q	Particularly fish species of the families: Scombridae, Clupeidae, Engraulidae, Coryfenidae, Pomatomidae, Scombresosidae.								
r	Single samples may be taken at retail level. In such a case the presumption laid down in Article 14(6) of Regulation (EC) No 178/2002, according to which the whole batch is to be deemed unsafe, shall not apply.								

References: 1. Malle P., Valle M., Bouquelet S. Assay of biogenic amines involved in fish decomposition. J. AOAC Internat. 1996, 79, 43-49. 2. Duflos G., Dervin C., Malle P., Bouquelet S. Relevance of matrix effect in determination s

of biogenic amines in plaice (*Pleuronectes platessa*) and whiting (*Merlangus merlangus*. J. AOAC Internat. 1999, 82, 1097-1101.

Interpretation of the test results

The limits given refer to each sample unit tested, excluding live bivalve molluscs and live echinoderms, tunicates and gastropods in relation to testing *E. coli*, where the limit refers to a pooled sample.

The test results demonstrate the microbiological quality of the batch tested⁽³⁾.

L. monocytogenes in ready-to-eat foods intended for infants and for special medical purposes:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

L. monocytogenes in ready-to-eat foods able to support the growth of *L. monocytogenes* before the food has left the immediate control of the producing food business operator when he is not able to demonstrate that the product will not exceed the limit of 100 cfu/g throughout the shelf-life:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

L. monocytogenes in other ready-to-eat foods and E. coli in live bivalve molluscs:

- satisfactory, if all the values observed are \leq the limit,
- unsatisfactory, if any of the values are > the limit.

Salmonella in different food categories:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

Staphylococcal enterotoxins in dairy products:

- satisfactory, if in all the sample units the enterotoxins are not detected,
- unsatisfactory, if the enterotoxins are detected in any of the sample units.

Enterobacter sakazakii in dried infant formulae and dried dietary foods for special medical purposes intended for infants below 6 months of age:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

Histamine in fishery products from fish species associated with a high amount of histidine:

- satisfactory, if the following requirements are fulfilled:
 - 1. the mean value observed is $\leq m$
 - 2. a maximum of c/n values observed are between m and M
 - 3. no values observed exceed the limit of M,
- unsatisfactory, if the mean value observed exceeds m or more than c/n values are between m and M or one or more of the values observed are > M.

Chapter 2.

Process hygiene criteria

2.1 *Meat and products thereof*

Food	Micro-	Samplin	g plan ^a	Limits ^b		Analytic		Action	
category	organisn	nsh	c	m	Μ	referenc method [°]	the	in case of unsatisfactory results	
2.1.1	Aerobic Carcases ofount cattle, sheep, goats and horses ^d			3,5 log cfu/cm ² daily mean log	5,0 log cfu/cm ² daily mean log	ISO 4833	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls	
	Enterobac	teriaceae		1,5 log cfu/cm ² daily mean log	2,5 log cfu/cm ² daily mean log	ISO 21528-2	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls	
2.1.2	Aerobic Carcones of count pigs			4,0 log cfu/cm ² daily mean log	5,0 log cfu/cm ² daily mean log	ISO 4833	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls	
	Enterobac	teriaceae		2,0 log cfu/cm ² daily mean log	3,0 log cfu/cm ² daily mean log	ISO 21528-2	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls	
2.1.3	<i>Salmonell</i> Carcases of cattle,	¢€0°	2 ^f	Absence area teste carcase		EN/ISO 6579	Carcases after dressing but	Improvements in slaughter hygiene,	

	sheep, goats and horses						before chilling	review of process controls and of origin of animals
2.1.4	Salmonel. Carcases of pigs	(&0°	5 ^f	Absence i area teste carcase		EN/ISO 6579	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls, origin of animals and of the biosecurity measures in the farms of origin
2.1.5	Salmonel. Poultry carcases of broilers and turkeys	[Φ0°	7 ^f	Absence in 25 g of a pooled sample of neck skin		EN/ISO 6579	Carcases after chilling	Improvements in slaughter hygiene and review of process controls, origin of animals and biosecurity measures in the farms of origin
2.1.6	Aerobic Winsfid meat count [#]	5	2	$5\times 10^5 \ cfu/g$	$5\times 10^6 \ cfu/g$	ISO 4833	End of the manufacto process	Improvements in upingluction hygiene and improvements in selection and/or

		5	2	50 of 1/2	500 sfr/	150	End	origin of raw materials
	E. coli ^h	5	2	50 cfu/g	500 cfu/ g	ISO 16649-1 or 2	End of the manufactu process	Improvements in upingluction hygiene and improvements in selection and/or origin of raw materials
2.1.7	Aerobic Mechanica separated meat (MSM) ⁱ	5 illy	2	5 × 10 ⁵ cfu/g	5 × 10 ⁶ cfu/g	ISO 4833	End of the manufactu process	Improvements in upingluction hygiene and improvements in selection and/or origin of raw materials
	E. coli ^h	5	2	50 cfu/g	500 cfu/ g	ISO 16649-1 or 2	End of the manufacto process	Improvements in upingluction hygiene and improvements in selection and/or origin of raw materials
2.1.8	<i>E. coli</i> ^h Meat preparation	5 ns	2	500 cfu/ g or cm ²	5 000 cfu/g or cm ²	ISO 16649-1 or 2	End of the manufactu process	Improvements in upingluction hygiene and improvements in selection and/or origin

Status: Point in time view as at 05/12/2007. Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 1441/2007. (See end of Document for details) of raw materials n = number of units comprising the sample; c = number of sample units giving values between m and M. a b For points 2.1.3-2.1.5 m = M. The most recent edition of the standard shall be used. с d The limits (m and M) shall apply only to samples taken by the destructive method. The daily mean log shall be calculated by first taking a log value of each individual test result and then calculating the mean of these log values. The 50 samples shall be derived from 10 consecutive sampling sessions in accordance with the sampling rules and frequencies laid down in this Regulation. f The number of samples where the presence of salmonella is detected. The c value is subject to review in order to take into account the progress made in reducing the salmonella prevalence. Member States or regions having low salmonella prevalence may use lower c values even before the review. This criterion shall not apply to minced meat produced at retail level when the shelf-life of the product is less then 24 g hours h E. coli is used here as an indicator of faecal contamination.

i These criteria apply to mechanically separated meat (MSM) produced with the techniques referred to in paragraph 3 of Chapter III of Section V of Annex III to Regulation (EC) No 853/2004 of the European Parliament and of the Council.

Interpretation of the test results

The limits given refer to each sample unit tested, excluding testing of carcases where the limits refer to pooled samples.

The test results demonstrate the microbiological quality of the process tested.

Enterobacteriaceae and aerobic colony count in carcases of cattle, sheep, goats, horses and pigs:

- satisfactory, if the daily mean log is \leq m,
- acceptable, if the daily mean log is between m and M,
- unsatisfactory, if the daily mean $\log i s > M$.

Salmonella in carcases:

- satisfactory, if the presence of *Salmonella* is detected in a maximum of c/n samples,
- unsatisfactory, if the presence of *Salmonella* is detected in more than c/n samples.

After each sampling session, the results of the last ten sampling sessions shall be assessed in order to obtain the n number of samples.

E. coli and aerobic colony count in minced meat, meat preparations and mechanically separated meat (MSM):

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.
- 2.2 *Milk and dairy products*

Food		Samplin	g plan ^a	Limits ^b		Analytic	0	Action
category	organisr	nsh	c	m	Μ	referenc method ^c	the	in case of unsatisfactory results

2.2.1	Enterobacteriaceae Pasteurised milk and other pasteurised liquid dairy products ^d	2	< 1/ml	5/ml	ISO 21528-1	End of the manufact process	Check on the ueffig: iency of heat treatment and prevention of recontamination as well as the quality of raw materials
2.2.2	<i>E. coli</i> ^e 5 Cheeses made from milk or whey that has undergone heat treatment	2	100 cfu/ g	1 000 cfu/g	ISO 16649-1 or 2	At the time during the manufact process when the <i>E</i> . <i>coli</i> count is expected to be highest ^f	Improvements in production hygiene using selection of raw materials
2.2.3	Coagulase5 Chosine Babbylococci from raw milk	2	10 ⁴ cfu/ g	10 ⁵ cfu/ g	EN/ISO 6888-2	At the time during the manufactu process when the number of staphyloc is expected to be highest	selection of raw materials. If values $> 10^5$ cfu/
2.2.4	Coagulase5 Positive Mathylococci from milk that has undergone a lower heat treatment than pasteurisation ^g and ripened cheeses made	2	100 cfu/ g	1 000 cfu/g	EN/ISO 6888-1 or 2		

		nges to legisla		e currently no	known outstan	ding effects fo cument for deta		
2.2.5	from milk or whey that has undergone pasteurisat or a stronger heat treatment ^g Coagulas, Untripped Staphyloc cheeses (fresh cheeses) made from milk or whey that has undergone pasteurisat or a stronger heat treatment ^g	ion e5 occi	2	10 cfu/g	100 cfu/ g	EN/ISO 6888-1 or 2	End of the manufact process	Improvements in upingluction hygiene. If values $> 10^5$ cfu/g are detected, the cheese batch has to be tested for staphylococca enterotoxins.
2.2.6	<i>E. coli</i> ^e Butter and cream made from raw milk or milk that has undergone a lower heat treatment than pasteurisat		2	10 cfu/g	100 cfu/ g	ISO 16649-1 or 2	End of the manufact process	Improvements in apingluction hygiene and selection of raw materials

2.2.7	Enterobact&riaceae Milk powder and whey powder ^d	0	10 cfu/g		ISO 21528-2	End of the manufactu process	Check on the refific iency of heat treatment and prevention of recontamination
	Coagulase5 positive staphylococci	2	10 cfu/g	100 cfu/ g	EN/ISO 6888-1 or 2	End of the manufactu process	Improvements in upingluction hygiene. If values $> 10^5$ cfu/g are detected, the batch has to be tested for staphylococcal enterotoxins.
2.2.8	Enterobact&riaceae Ice cream ^h and frozen dairy desserts	2	10 cfu/g	100 cfu/ g	ISO 21528-2	End of the manufactu process	Improvements in upingluction hygiene
2.2.9	Enterobacteoiaceae Dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age	0	Absence	in 10 g	ISO 21528-1	End of the manufactu process	Improvements in upingluction hygiene to minimise contamination ⁱ

2.2	.10	Enterobac Dried follow- on formulae	teriaceae	0	Absence	in 10 g	ISO 21528-1	End of the manufactu process	Improvements in apingluction hygiene to minimise contamination
2.2		Presumpt Detectilus infant _{us} formulae and dried dietary foods for special medical purposes intended for infants below six months of age	ive	1	50 cfu/g	500 cfu/ g	EN/ISO 7932 ⁱ	End of the manufactu process	Improvements in apingluction hygiene. Prevention of recontamination Selection of raw material.
a	n = nun	nber of units co	omprising the s	ample; c = nu	nber of sampl	e units giving	values betweer	n m and M.	
b	For poi	nts 2.2.7, 2.2.9	and 2.2.10 m	= M.					
c	The mo	st recent edition	on of the standa	ard shall be use	ed.				
d	The cri	terion shall not	apply to produ	ucts intended f	or further proc	essing in the f	ood industry.		
e	E. coli	is used here as	an indicator fo	or the level of h	iygiene.				
f	For cheeses which are not able to support the growth of E. coli, the E. coli count is usually the highest at the beginning of the ripening period, and for cheeses which are able to support the growth of E. coli, it is normally at the end of the ripening period.								
g		ing cheeses wh does not pose				ne satisfaction	of the compete	nt authorities,	that the
h	Only ice creams containing milk ingredients.								
i	organis samples to demo	ms has been es s tested in such	tablished at an a plant, the ba	individual pla ttch has to be t	nt level. If En ested for E. sa	terobacteriacea kazakii. It shal	e are detected l be the respon	n between these in any of the p sibility of the r between Enter	roduct nanufacturer
	1 1 6	in coulum is n	ated on a Petri	dish of 140 m	m diameter or	on three Detri	dishes of 90 m	um diameter	

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

Enterobacteriaceae in dried infant formulae, dried dietary foods for special medical purposes intended for infants below six months of age and dried follow-on formulae:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

E. coli, Enterobacteriaceae (other food categories) and coagulase-positive staphylococci:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are $\leq m$,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Presumptive Bacillus cereus in dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are $\leq m$,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Food	Micro-	Samplin	g planª	Limits	Limits		aStage	Action
category	organisn	ns _h	c	m	Μ	referenc method ^b	_	in case of unsatisfactory results
	Enterobac Egg products	teriaceae	2	10 cfu/g or ml	100 cfu/ g or ml	ISO 21528-2	End of the manufactu process	Checks on the reiffig ciency of the heat treatment and prevention of recontaminatior

2.3 Egg products

n = number of units comprising the sample; c = number of sample units giving values between m and M.

b The most recent edition of the standard shall be used.

Interpretation of the test results

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

Enterobacteriaceae in egg products:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are $\leq m$,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

2.4 *Fishery products*

Food category	Micro-	Samplin	g plan ^a	Limits		Analytic		Action
	organisr	ns _h	c	m	М	referenc method ^b		in case of unsatisfactory results
2.4.1	<i>E. coli</i> Shelled and shucked products	5	2	1/g	10/g	ISO TS 16649-3	End of the manufactu process	Improvements in apingluction hygiene
	of oagulase cposited/e crtsphydaa and molluscan shellfish	occi	2	100 cfu/ g	1 000 cfu/g	EN/ISO 6888-1 or 2	End of the manufactu process	Improvements in upingluction hygiene
a n = nur	nber of units co	omprising the	sample; c = nu	mber of sampl	e units giving	values between	m and M.	
b The mo	ost recent edition	on of the stand	ard shall be use	ed.				

Interpretation of the test results

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

E. coli in shelled and shucked products of cooked crustaceans and molluscan shellfish:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Coagulase-positive staphylococci in shelled and cooked crustaceans and molluscan shellfish:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

	Food category	Micro- organism	Sampling plan ^a		Limits		AnalyticaB		Action	
ca			nsh	c	m	Μ	method ^b the ot criterion u	in case of unsatisfact results	of unsatisfactory	
a	a $n =$ number of units comprising the sample; $c =$ number of sample units giving values between m and M.									
b	The mos	st recent editio	n of the standa	ard shall be use	:d.					

2.5 *Vegetables, fruits and products thereof*

2.5.1	<i>E. coli</i> Precut fruit and vegetables (ready- to- eat)	5	2	100 cfu/ g	1 000 cfu/g	ISO 16649-1 or 2	Manufact process	in production hygiene, selection of raw materials
2.5.2	<i>E. coli</i> Unpasteuri fruit and vegetable juices (ready- to- eat)	5 sed	2	100 cfu/ g	1 000 cfu/g	ISO 16649-1 or 2	Manufact process	utimgrovements in production hygiene, selection of raw materials

b The most recent edition of the standard shall be used.

Interpretation of the test results

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

E. coli in precut fruit and vegetables (ready-to-eat) and in unpasteurised fruit and vegetable juices (ready-to-eat):

- satisfactory, if all the values observed are \leq m,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Chapter 3.

Rules for sampling and preparation of test samples

3.1 *General rules for sampling and preparation of test samples*

In the absence of more specific rules on sampling and preparation of test samples, the relevant standards of the ISO (International Organisation for Standardisation) and the guidelines of the Codex Alimentarius shall be used as reference methods.

3.2 Bacteriological sampling in slaughterhouses and at premises producing minced meat and meat preparations

Sampling rules for carcases of cattle, pigs, sheep, goats and horses

The destructive and non-destructive sampling methods, the selection of the sampling sites and the rules for storage and transport of samples are described in standard ISO 17604.

Five carcases shall be sampled at random during each sampling session. Sample sites must be selected taking into account the slaughter technology used in each plant.

When sampling for analyses of Enterobacteriaceae and aerobic colony counts, four sites of each carcase shall be sampled. Four tissue samples representing a total of 20 cm^2 shall be obtained by the destructive method. When using the non-destructive method for this purpose, the sampling area shall cover a minimum of 100 cm^2 (50 cm² for small ruminant carcases) per sampling site.

When sampling for *Salmonella* analyses, an abrasive sponge sampling method shall be used. Areas most likely to be contaminated shall be selected. The total sampling area shall cover a minimum of 400 cm^2 .

When samples are taken from the different sampling sites on the carcase, they shall be pooled before examination.

Sampling rules for poultry carcases

For the Salmonella analyses, a minimum of 15 carcases shall be sampled at random during each sampling session and after chilling. A piece of approximately 10 g from neck skin shall be obtained from each carcase. On each occasion the neck skin samples from three carcases shall be pooled before examination in order to form 5×25 g final samples. *Guidelines for sampling*

More detailed guidelines on the sampling of carcases, in particular concerning the sampling sites, may be included in the guides to good practice referred to in Article 7 of Regulation (EC) No 852/2004.

Sampling frequencies for carcases, minced meat, meat preparations and mechanically separated meat

The food business operators of slaughterhouses or establishments producing minced meat, meat preparations or mechanically separated meat shall take samples for microbiological analysis at least once a week. The day of sampling shall be changed each week to ensure that each day of the week is covered.

As regards the sampling of minced meat and meat preparations for *E. coli* and aerobic colony count analyses and the sampling of carcases for Enterobacteriaceae and aerobic colony count analyses, the frequency may be reduced to fortnightly testing if satisfactory results are obtained for six consecutive weeks.

In the case of sampling for *Salmonella* analyses of minced meat, meat preparations and carcases, the frequency may be reduced to fortnightly if satisfactory results have been obtained for 30 consecutive weeks. The salmonella sampling frequency may also be reduced if there is a national or regional salmonella control programme in place and if this programme includes testing that replaces the sampling described in this paragraph. The sampling frequency may be further reduced if the national or regional salmonella control programme demonstrates that the salmonella prevalence is low in animals purchased by the slaughterhouse.

However, when justified on the basis of a risk analysis and consequently authorised by the competent authority, small slaughterhouses and establishments producing minced meat and meat preparations in small quantities may be exempted from these sampling frequencies.

- (1) OJ L 139, 30.4.2004, p. 1, as corrected by OJ L 226, 25.6.2004, p. 3.
- (2) OJ L 338, 22.12.2005, p. 1.
- (3) The test results may be used also for demonstrating the effectiveness of the hazard analysis and critical control point principles or good hygiene procedure of the process.

Status:

Point in time view as at 05/12/2007.

Changes to legislation:

There are currently no known outstanding effects for the Commission Regulation (EC) No 1441/2007.