

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

Green tea (Camellia sinensis) leaves and stalks, fermented or otherwise, Camellia sinensis)	Dibromoethane Oxide (HDBO)	Metiram	T
		Propineb	
		Zineb	

6. TEA

leaves of
and alpha
stalks, and
fermented beta
or
otherwise,
Camellia
sinensis)

7. HOPS (dried)

hop pellets & unconcentrated powder

Green tea (Camellia sinensis) leaves and stalks, fermented or otherwise, Camellia sinensis)	Dibromoethane Oxide (HDBO)	Metiram	T
		Propineb	
		Zineb	

8. CEREALS

Wheat of alpha & beta

Rye of alpha & beta

Bamboo of alpha & beta

Sorghum of alpha

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

Chemical	Meat ⁽³⁾	Milk & Dairy produce ⁽⁵⁾	Eggs
Chlorpyrifos	0.01*	0.01*	0.01*
Disulfoton	0.01*	0.01*	0.01*
Endosulfan	0.01*	0.01*	0.01*
Phosphamidon	0.01*	0.01*	0.01*
Permethrin	0.01*	0.01*	0.01*
Triphenylethylene pyrethroids	0.01*	0.01*	0.01*
Dibromethane	0.02	0.02	0.02
βγ			
Metiram	0.02*	0.02*	0.02*
Propineb	0.02*	0.02*	0.02*
Zinc	0.02*	0.02*	0.02*

FOOTNOTES

UNITS:

Maximum residue levels (MRLs) are expressed in milligrammes of residue per kilogramme of food.

KEY:

* Level at or about the limit of determination.

FOOTNOTES:

1. Paddy or rough rice, husked rice and semi-milled or wholly milled rice.
2. Other cereals do not include rice.
3. Levels are measured on fat, except in the case of foods with a fat content of 10% or less by weight. In these cases the residue is related to the total weight of the boned foodstuff and the MRL is one tenth of the value given in the table, but must be no less than 0.01 mg/kg.
4. These levels are for fresh raw cow’s milk and fresh whole cream cow’s milk expressed on the whole milk.
5. For preserved, concentrated or sweetened cow’s milk; for raw milk and whole cream milk of another origin; and for butter, cheese or curd whether made from cow’s milk or other milk of a combination, the following levels apply:
 - if the fat content is less than 2% by weight. the MRL is taken as half that set for raw milk and whole cream milk;
 - if the content is 2% or more by weight, the MRL is expressed in mg/kg of fat and is set at 25 times that set for raw milk and whole cream milk.
6. Birds' eggs in shell (other than eggs for hatching) and whole egg products and egg yolk products (whether fresh, dried or otherwise prepared).
7. Sheepmeat only.
8. Poultrymeat only.
9. All meat except sheepmeat.
10. Other meat products.
11. All meat except poultrymeat.
12. Pig kidney.

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

13. Cattle, goat and sheep kidney.
14. Procymidone:
 - 1 mg/kg applies to whole seed
 - 0.05 mg/kg applies to seed without shell.
15. All meat except liver and kidney.
16. Ruminant liver.
17. All meat except ruminant liver.
18. For animal products MRLs relate to cyhalothrin (sum of isomers)
19. With the exception of meat and other ovine, bovine and caprine products.
20. Footnotes 3, 5 and 6 do not apply in cases where the lower limit of analytical determination is indicated.
21. Meat of cattle, sheep and goats.
22. Other than meat or liver of cattle, sheep and goats, and poultry meat.
23. Liver of cattle, sheep and goats. The residue definition for this MRL is: 1,1-bis-(parachlorophenol)-2,2-dichloroethanol (PP'-FW152), expressed as dicofol.
24. Fat, liver and kidney.
25. Other than fat, liver and kidney.
26. The residues definition for these MRLs is: sum of propyzamide and all metabolites containing the 3,5-dichlorobenzoic acid fraction expressed as propyzime.