

## ANNEX II

## MICRO-ORGANISMS

Number (or EC number)	Additive	Chemical formula, or description	Species or category of animal	Maximum age	Minimum	Maximum	Other provisions	End of period of authorisation
					content	content		
					CFU/ kg of complete feedingstuff	CFU/ kg of complete feedingstuff		
1	<i>Bacillus cereus</i> var. <i>toyoi</i> NCIMB 40112/ CNCM I-1012	Preparation of <i>Bacillus cereus</i> var. <i>toyoi</i> containing a minimum of $1 \times 10^{10}$ CFU/g additive	Chickens for fattening	—	$0,2 \times 10^9$	$1 \times 10^9$	In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability to pelleting May be used in compound feed containing the permitted coccidiostats: monensin sodium, lasalocid sodium, salinomycin sodium, decoquinate, robenidine, narasin, halofuginone	7.10.2004
			Laying hens	—	$0,2 \times 10^9$	$1 \times 10^9$	In the directions for use of the additive and premixture, indicate	7.10.2004

**Changes to legislation:** There are currently no known outstanding effects for the Commission Regulation (EC) No 256/2002, ANNEX II. (See end of Document for details)

				the storage temperature, storage life and stability to pelleting	
Calves	6 months	$0,5 \times 10^9$	$1 \times 10^9$	In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability to pelleting	7.10.2004
Cattle for fattening	—	$0,2 \times 10^9$	$0,2 \times 10^9$	In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability to pelleting The quantity of <i>Bacillus cereus</i> var. <i>toyoi</i> in the daily ration must not exceed $1,0 \times 10^9$	7.10.2004

**Changes to legislation:** There are currently no known outstanding effects for the Commission Regulation (EC) No 256/2002, ANNEX II. (See end of Document for details)

						CFU for 100 kg body weight. Add $0,2 \times 10^9$ CFU for each additional 100 kg body weight	
		Breeding does	—	$0,1 \times 10^9$	$5 \times 10^9$	In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability to pelleting May be used in compound feed containing the permitted coccidiostat: robenidine	7.10.2004
		Rabbits for fattening	—	$0,1 \times 10^9$	$5 \times 10^9$	In the directions for use of the additive and premixture, indicate the storage temperature, storage life and stability	7.10.2004

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 256/2002, ANNEX II. (See end of Document for details)*

---

									to pelleting May be used in compound feed containing the permitted coccidiostats: robenidine, salinomycin sodium	
--	--	--	--	--	--	--	--	--	--	--

---

**Changes to legislation:**

There are currently no known outstanding effects for the Commission Regulation (EC) No 256/2002, ANNEX II.