

ANNEX I

PART A

General provisions

1. Where there is more than one group of nutritional characteristics indicated in column 2 of Part B, denoted by ‘and/or’, for the same nutritional purpose, the manufacturer has the option to use either or both groups of essential characteristics, in order to achieve the nutritional purpose defined in column 1. For each option the corresponding labelling declarations are given opposite in column 4.
2. Where a group of additives is mentioned in column 2 or column 4 of Part B the additive(s) used must be authorised in Regulation (EC) No 1831/2003 of the European Parliament and of the Council⁽¹⁾ as corresponding to the specified essential characteristic.
3. Where the source(s) of ingredients or of analytical constituents is (are) required in column 4 of Part B, the manufacturer must make a precise declaration (e.g. specific name of the ingredient(s), animal species or part of the animal) allowing the evaluation of conformity of the feedingstuff with the corresponding essential nutritional characteristics.
4. Where the declaration of a substance, also authorised as an additive, is required in column 4 of Part B and is accompanied by the expression ‘total’ the declared content must refer to, as appropriate, the quantity naturally present where none is added or, by derogation from Council Directive 70/524/EEC⁽²⁾, the total quantity of the substance naturally present and the amount added as an additive.
5. The declarations required in column 4 of Part B with the reference ‘if added’ are compulsory where the ingredient or the additive has been incorporated or increased specifically to enable the achievement of the particular nutritional purpose.
6. The declarations to be given in accordance with column 4 of Part B concerning analytical constituents and additives must be quantitative.
7. The recommended period of use indicated in column 5 of part B indicates a range within which the nutritional purpose should normally be achieved. Manufacturers can refer to more precise periods of use, within the fixed limits.
8. Where a feedingstuff is intended to meet more than one particular nutritional purpose, it must comply with the corresponding entries in Part B.
9. In the case of complementary feedingstuffs intended for particular nutritional purposes guidance on the balance of the daily ration must be provided in the instructions for use contained on the label.
- [^{F1}10. When a feed intended for particular nutritional purposes is placed on the market in the form of a bolus, being a feed material or complementary feed intended for individual oral administration by forced feeding, the label of the feed shall, if applicable, mention the maximum period of continuously release of the bolus and the daily release rate for each additive for which a maximum content in complete feed is fixed. At the request of the competent authority the feed business operator which is placing a bolus on the market shall provide proof that the daily available additive level in the digestive tract will not exceed, if applicable, the maximum content of the additive established

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

per kg complete feed during the whole feeding period (slow release effect). It is recommended that feed in form of bolus is administered by a veterinarian or any other competent person.]

Textual Amendments

- F1** Inserted by [Commission Regulation \(EU\) No 5/2014 of 6 January 2014 amending Directive 2008/38/EC establishing a list of intended uses of animal feedingstuffs for particular nutritional purposes \(Text with EEA relevance\)](#).

PART B

List of intended uses

Particular nutritional purpose	Essential nutritional characteristics	Species or category of animals	Labelling declarations	Recommended length of time	Other provisions
(1)	(2)	(3)	(4)	(5)	(6)
[^{F2} Support of renal function in case of chronic renal insufficiency ^a	High quality proteins and restricted levels of phosphorus maximum 5 g/kg complete pet food with a moisture content of 12 % and crude protein maximum 220 g/kg complete pet food with a moisture content of 12 % or	Dogs	— Protein source(s) — Calcium — Phosphorus — Potassium — Sodium — Content of essential fatty acids (if added)	Initially up to 6 months ^b	— The feed shall be placed on the market as complete feed. — Recommended digestibility of proteins: minimum 85 %. — Indicate on the labelling: ‘It is recommended that advice from a veterinarian be sought before

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

				available at all times.'
High quality proteins and restricted levels of phosphorus maximum 6,2 g/kg complete pet food with a moisture content of 12 % and crude protein maximum 320 g/kg complete pet food with a moisture content of 12 % or	Cats	— — — — —	Protein source(s) Calcium Phosphorus Potassium Sodium Content of essential fatty acids (if added)	Initially up to 6 months ^b — The feed shall be placed on the market as complete feed. — Recommended digestibility of proteins: minimum 85 %. — Indicate on the labelling: 'It is recommended that advice from a veterinarian be sought before use and before extending the period of use.' — Indicate in the instructions for use: 'Water should

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

	protein but of high quality				It is recommended that a veterinarian's opinion be sought before use.
	— Urine acidifying properties and low level of magnesium	Cats	— — — — — — — —	Calcium Phosphorus Sodium Magnesium Potassium Chlorides Sulphur Total taurine Urine acidifying substances	
Reduction of struvite stone recurrence ^c	Urine acidifying properties and moderate level of magnesium	Dogs and cats	— — — — — — —	Calcium Phosphorus Sodium Magnesium Potassium Chlorides Sulphur Urine acidifying substances	Up to 6 months Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use.
Reduction of urate stones formation	Low level of purines, low level of protein but of high quality	Dogs and cats	Protein source(s)	Up to 6 months but lifetime use in cases of irreversible disturbance of uric acid metabolism	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use.
Reduction of oxalate stones formation	Low level of calcium, low level of vitamin D, and urine alkalisating properties	Dogs and cats	— — — — — — — —	Phosphorus Calcium Sodium Magnesium Potassium Chlorides Sulphur Total vitamin D Hydroxyproline	Up to 6 months Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use.

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

			—	Urine alkalisng substances	
Reduction of cystine stones formation	Low level of protein, moderate level of sulphur amino acids and urine alkalisng properties	Dogs and cats	— — — — —	Total sulphur amino acids Sodium Potassium Chlorides Sulphur Urine alkalisng substances	Initially up to 1 year Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use.
Reduction of ingredient and nutrient intolerances ^d	— Selected protein source(s)	Dogs and cats	— —	Protein source(s) Content of essential fatty acids (if added)	if to 8 weeks: signs of intolerance disappear this feed can be used indefinitely
	and/or — Selected carbohydrate source(s)		— —	Carbohydrate source(s) Content of essential fatty acids (if added)	
Reduction of acute intestinal absorptive disorders	Increased level of electrolytes and highly digestible ingredients	Dogs and cats	— — —	Highly digestible ingredients including their treatment if appropriate Sodium Potassium Source(s) of mucilaginous	Indicate on the package, container or label: — During periods of and recovery from acute diarrhoea.

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

			substances (if added)	—	It is recommended that a veterinarian's opinion be sought before use.
Compensation for maldigestion ^e	Highly digestible ingredients and low level of fat	Dogs and cats	Highly digestible ingredients including their treatment if appropriate	3 to 12 weeks, but lifetime in the case of chronic pancreatic insufficiency	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use.
[^{F2} Support of heart function in the case of chronic cardiac insufficiency	Level of sodium restricted to be lower than 2,6 g/kg complete pet food with a moisture content of 12 %	Dogs and cats	— Magnesium — Potassium — Sodium	Initially up to 6 months	— The feed shall be placed on the market as complete feed. — Indicate on the labelling: 'It is recommended that advice from a veterinarian be sought before use and before extending the period

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					of use.']
Regulation of glucose supply (<i>Diabetes mellitus</i>)	Low level of rapid glucose-releasing carbohydrates	Dogs and cats	— — — — — — —	Carbohydrates initially up to 6 months Treatment of carbohydrates if appropriate Starch Total sugar Fructose (if added) Content of essential fatty acids (if added) Source(s) of short and medium chain fatty acids (if added)	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use.
Support of liver function in the case of chronic liver insufficiency	— High quality protein, moderate level of protein, high level of essential fatty acids and high level of highly	Dogs	— — — —	Protein initially up to 6 months Content of essential fatty acids Highly digestible carbohydrates including their treatment if appropriate Sodium Total copper	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use. Indicate in the instructions for use: Water should be

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

		digestible carbohydrates				available at all times.
	—	Cats High quality protein, moderate level of protein and high level of essential fatty acids	— — — —	Protein source Content of essential fatty acids Sodium Total copper	Initially up to 6 months	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use. Indicate in the instructions for use: Water should be available at all times.
Regulation of lipid metabolism in the case of hyperlipidaemia	Low level of fat and high level of essential fatty acids	Dogs and cats	— —	Content of essential fatty acids Content of n-3 fatty acids (if added)	Initially up to 2 months	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use.
[F2] Reduction of iodine levels in feed in case of hyperthyroidism	Restricted level of iodine: maximum 0,26 mg/kg complete pet food with a moisture content of 12 %	Cats	Total iodine		Initially up to 3 months	— The feed shall be placed on the market as complete feed.

					—	Indicate on the labelling: 'It is recommended that advice from a veterinarian be sought before use and before extending the period of use.'
[^{F2} Reduction of copper in the liver	Restricted level of copper: maximum 8,8 mg/kg complete pet food with a moisture content of 12 %	Dogs	Total copper	Initially up to 6 months	—	The feed shall be placed on the market as complete feed. Indicate on the labelling: 'It is recommended that advice from a veterinarian be sought before use and before extending the

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					period of use.']
Reduction of excessive body weight	Low energy density	Dogs and cats	Energy value	Until target body weight is achieved	In the instructions for use an appropriate daily intake must be recommended.
[^{F3} Nutritional restoration, convalescence]	High energy density, high concentrations of essential nutrients and highly digestible ingredients	Dogs and cats	— Highly digestible ingredients including their treatment if appropriate — Energy value — Content of n-3 and n-6 fatty acids (if added)	Until restoration is achieved	In the case of feed specially presented to be given via tubing, indicate on the labelling: 'Administration under veterinary supervision.'
[^{F2}	The complementary feed may contain <i>Enterococcus faecium</i> DSM 10663/ NCIMB 10415 in a concentration higher than 100 times the relevant fixed maximum content in complete feed.	Dogs	<i>Enterococcus faecium</i> DSM 10663/ NCIMB 10415 including added amount	10 to 15 days	— The instructions for use of the feed shall ensure that the legal maximum content of the gut flora stabiliser// micro-organism for

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					complete feed is respected. Indicate on the labelling: 'It is recommended that advice from a veterinarian be sought before use and before extending the period of use.']]
Support of skin function in the case of dermatosis and excessive loss of hair	High level of essential fatty acids	Dogs and cats	Content of essential fatty acids	Up to 2 months	Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use.
[^{F5} Support of the metabolism of joints in the case of osteoarthritis	Dogs: Minimum dry matter content of total omega-3 fatty acids 3,3 % and of eicosapentaenoic acid	Dogs and cats	Dogs: — Total omega-3 fatty acids — Total EPA — Total vitamin E Cats: — Total omega-3 fatty acids	Initially up to 3 months	It is recommended that a veterinary surgeon's opinion be sought before use or before extending the period of use.]

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

		(EPA) 0,38 %. Adequate level of vitamin E. Cats: Minimum dry matter content of total omega-3 fatty acids 1,2 % and of docosahexaenoic acid (DHA) 0,28 %. Increased levels of methionine and manganese. Adequate level of vitamin E.	— — — — —	Total DHA Total methionine Total manganese Total vitamin E	
Reduction of the risk of milk fever	—	Low level of calcium Dairy cows	— — —	Calcium to 4 weeks Phosphorus Magnesium	Indicate in the instructions for use: Stop feeding after calving
	and/or				
	—	Low cations/anions ratio	— — — — —	Calcium to 4 weeks Phosphorus Sodium Potassium Chlorides Sulphur	Indicate in the instructions for use: Stop feeding after calving

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

or				
—	High level of zeolite (synthetic sodium aluminium silicate)	Content of synthetic sodium aluminium silicate	The 2 weeks before calving	Indicate in the instructions for use: — The amount of feed shall be restricted to ensure that a daily intake of 500 g sodium aluminium-silicate per animal is not exceeded. — Stop feeding after calving
or				
—	High level of calcium in the form of highly available calcium salts	Total calcium content, sources and respective quantity of calcium	Star at first signs of parturition to two days subsequent to parturition	Indicate on the package, container or label: — The instruction of use i.e. the number of applications and the time before and

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					— after calving. The text 'It is recommended that a nutritional expert's opinion be sought before use'
Reduction of the risk of ketosis ^{gh}	Ingredients providing glucogenic energy sources	Dairy cows and ewes	— — —	Ingredients providing glucogenic energy sources before and after calving ⁱ Propanediol (if added as a glucose precursor) Glycerol (if added as a glucose precursor)	6 weeks before and the first 3 weeks after lambing ^j
Reduction of the risk of tetany (hypomagnesaemia)	High level of magnesium, easily available carbohydrates, moderate level of protein and low level of potassium	Ruminants	— — — —	Starch Total sugars Magnesium Sodium Potassium	3 to 10 weeks during fasting periods of grass growth
					In the instructions for use guidance shall be provided on the balance of the daily ration, with regard to the inclusion of fibre and easily available energy sources. In the case of feedingstuffs for ovines

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					indicate on the package, container or label: Especially for lactating ewes.	
Reduction of the risk of acidosis	Low level of easily fermentable carbohydrates and high buffering capacity	Ruminants	— —	Starch Total sugars	Maximum 2 months ^k	In the instructions for use guidance shall be provided on the balance of the daily ration, with regard to the inclusion of fibre and easily fermentable carbohydrate sources. In the case of feedingstuffs for dairy cows indicate on the package, container or label: Especially for high yielding cows. In the case of feedingstuffs for ruminants for fattening indicate on the package, container or label: Especially for intensively fed. ^l
[^{F1} Long-term supply of grazing	High level of — Trace with a element	Ruminants with a functional rumen	—	Name and total amounts	Up to 12 months	— Application in the form

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

<p>animals with trace elements and/or vitamins</p>	<p>—</p> <p>The complementary feed may contain feed additives in a concentration higher than 100 times the relevant fixed maximum content in complete feed.</p>	<p>and/or vitamins, provitamins and chemically well-defined substances having similar effects.</p>	<p>—</p> <p>of each added trace element, vitamin, provitamin and chemically well-defined substance having similar effects. Daily release rate for each trace element and/or vitamin if a bolus is used; Maximum period of continuously release of the trace element or vitamin if a bolus is used.</p>	<p>of bolus is allowed. A bolus may contain up to 20 % iron in an inert, non-bioavailable form, in order to increase its density. Indicate on the labelling of the feed: — Simultaneous supplementation of additives with a maximum content from other sources to those incorporated in a bolus, if applicable, shall be avoided.</p>
--	---	--	--	---

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

Compensation for insufficient iron availability after birth	High level of iron compounds authorised under the functional group	Suckling piglets and calves	Total iron content	After birth up to 3 weeks	The instructions for use for the feed shall ensure that the legal maximum

— Before using, it is recommended to have advice from a veterinarian or nutritionist concerning:

(1) the balance of trace elements in the daily ration;

(2) the trace element status of the herd

— The bolus contains x % inert iron in order to increase its density, if applicable.

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

	‘compounds of trace elements’ of the category ‘nutritional additives’ as referred to in Annex I to Regulation (EC) No 1831/2003. The complementary feed may contain iron in a concentration higher than 100 times the relevant fixed maximum content in complete feed.					contents of iron for complete feed are respected.
Support the regeneration of hooves, trotters and skin	High level of zinc. The complementary feed may contain zinc in a concentration higher than 100 times the relevant fixed maximum content in complete feed.	Horses, ruminants and pigs	Total amount of — zinc — methionine	Up to 8 weeks		The instructions of use for the feed shall ensure that the legal maximum contents of zinc for complete feed are respected.
Support the preparation for oestrus and reproduction	— High level of selenium and a minimum content of vitamin E per kg	Mammals	Names and total amounts of each added trace element and vitamin.	— Cows: 2 weeks before the end of gestation until the next gestation	—	The instructions of use for the feed shall ensure that the respective legal

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

<p>—</p>	<p>complete feed with a moisture content of 12 % for pigs of 53 mg, rabbits of 35 mg, for dogs, cats, mink of 88 mg; a minimum content of vitamin E per animal and day for ovines of 100 mg, cattle of 300 mg, horses of 1 100 mg or High level(s) of vitamins A and/ or vitamin D and/ or a</p>	<p>—</p>	<p>is confirmed. Sows: 7 days before until 3 days after parturition and 7 days before until 3 days mating. Other female mammals: from the last part of gestation until the next gestation is confirmed. Males: during periods of reproductive activity.</p>	<p>maximum contents for complete feed are respected. Indicate on the labelling of the feed: ‘Guidance on the situations in which the use of the feed is appropriate.’</p>
----------	--	----------	---	---

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

	<p>minimum content of beta-carotene of 300 mg per animal and day.</p> <p>The complementary feed may contain selenium, vitamin A and D in a concentration higher than 100 times the relevant fixed maximum content in complete feed.</p>			
—	<p>High level(s) of vitamin A and/or vitamin D or High level(s) of selenium and/or zinc and/or a minimum content of vitamin E of 44 mg/kg</p>	Birds	Names and total amounts of each added trace element and vitamin.	<p>— For females: during oestrus</p> <p>— For males: during periods of reproductive activity]</p>

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

		complete feed with a moisture content of 12 %.				
		The complementary feed may contain selenium, zinc, vitamin A and D in a concentration higher than 100 times the relevant fixed maximum content in complete feed.				
[F2] Stabilisation of water and electrolyte balance to support the physiological digestion	—	Predominantly, pigs, kids and foals	—	Sodium to 7 days	—	Recommended range of electrolytes per litre portion ready prepared for feeding: Sodium: 1,7 g — 3,5 g Potassium: 0,4 g — 2,0 g Chlorides: 1 g — 2,8 g
	—	Electrolytes, sodium and potassium chlorides Buffering capacity ^m : minimum 60 mmol per litre portion ready prepared for feeding Easily digestible carbohydrates	—	Potassium Chlorides Carbohydrate source(s) Bicarbonates and/or citrates (if added)	—	

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

						(1)	'In case of risk of, during periods of, or recovery from digestive disturbance (diarrhoea).'
						(2)	'It is recommended that advice from a veterinarian be sought before use and before extending the period of use.'
						—	Indicate in the instructions for use:
						(1)	The recommended intake of the ready stirred potion and of milk, if appropriate.

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					(2) In case of bicarbonates and/or citrates are above 40 mmol per litre potion ready prepared for feeding: 'Simultaneous feeding of milk should be avoided in animals with abomasum']	
Reduction of the risk of urinary calculi	Low level of phosphorus, magnesium, and urine acidifying properties	Ruminants	— — — — — — —	Calcium Phosphorus Sodium Magnesium Potassium Chlorides Sulphur Urine acidifying substances	Up to 6 weeks	Indicate on the package, container or label: Especially for intensively fed young animals. Indicate in the instructions for use: Water should be available at all times.
Reduction of stress reactions	— High level magnesium and/or	Pigs	—	Magnesium	1 to 7 days	Guidance shall be provided on the situation in which the use of

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

	— Highly digestible ingredients		— Highly digestible ingredients including their treatment if appropriate — Content of n-3 fatty acids (if added)	this feed is appropriate.
[F3]Stabilisation of physiological digestion	Low buffering capacity, and highly digestible ingredients	Piglets	— Highly digestible ingredients including their treatment if appropriate — Buffering capacity — Source(s) of astringent substances (if added) — Source(s) of mucilaginous substances (if added)	Indicate on the labelling: 'In case of risk of, during periods of, or recovery from, digestive disturbance.'
	Highly digestible ingredients	Pigs	— Highly digestible ingredients including their treatment if appropriate — Source(s) of astringent substances (if added)	

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

			— Source(s) of mucilaginous substances (if added)		
[^{F2}	<p>Feed additives of the functional group ‘Gut flora stabiliser’ of the category ‘zotechnical additives’ as referred to in Annex I to Regulation (EC) No 1831/2003 or, pending the re-authorisation procedure as referred to in Article 10 of Regulation (EC) No 1831/2003, feed additives of the group ‘micro-organisms’. The complementary feed may contain feed additives of the functional group of ‘gut flora stabilisers/ micro-organisms’ in a concentration higher than 100 times the relevant fixed maximum content in</p>	<p>Animal species for which the gut flora stabiliser/ micro-organism is authorised</p>	<p>Name and added amount of the gut flora stabiliser/ micro-organism</p>	<p>Up to 4 weeks</p>	<p>— Indicate on the labelling of the feed:</p> <p>(1) ‘In case of risk of, during periods of, or recovery from digestive disturbance’.</p> <p>(2) If applicable: ‘The feed contains a gut flora stabiliser/ micro-organism in a concentration above 100 times the authorised maximum content in complete feed’.</p> <p>— The instructions for</p>

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

	complete feed.				use of the feed shall ensure that the legal maximum content of the gut flora stabiliser/ micro-organism for complete feed is respected.]]
Reduction of the risk of constipation	Ingredients stimulating intestinal passage	Sows	Ingredients stimulating intestinal passage	10 to 14 days before and 10 to 14 days after farrowing	
Reduction of the risk of fatty liver syndrome	Low energy and high proportion of metabolisable energy from lipids with high level of polyunsaturated fatty acids	Laying hens	— — —	Energy value (calculated according to EC method) Percentage of metabolisable energy from lipids Content of poly-unsaturated fatty acids	Up to 12 weeks
Compensation for malabsorption	Low level of saturated fatty acids and high level	Poultry excluding geese and pigeons	—	Percentage of saturated fatty	During the first 2 weeks after hatching

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

	of fat soluble vitamins		—	acids in relation to total fatty acids Total vitamin A Total vitamin D Total vitamin E Total vitamin K	
Compensation for chronic insufficiency of small intestine function	Highly precaecally digestible carbohydrates proteins and fats	Equines ^a	—	Source(s) of highly digestible carbohydrates, proteins and fats, including their treatment if appropriate	Initially up to six months Guidance should be provided on the situations in which the use of the feed is appropriate and the manner in which it should be fed including many small meals per day. Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use.

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

Compensation for chronic digestive disorders of large intestine	Highly digestible fibre	Equines	— —	Fibre source(s) Content of n-3 fatty acids (if added)	Initially up to six months	Guidance should be provided on the situations in which the use of the feed is appropriate and the manner in which the feed should be fed. Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use.
Reduction of stress reactions	Highly digestible ingredients	Equines	— — —	Magnesium Highly digestible ingredients including their treatment if appropriate Content of n-3 fatty acids (if added)	From two to four weeks	Guidance shall be provided on the precise situations in which the use of the feed is appropriate.
Compensation of electrolyte loss in the cases of heavy sweating	Predominantly electrolytes and easily absorbable carbohydrates	Equines	— — — — —	Calcium Sodium Magnesium Potassium Chlorides Glucose	One to three days	Guidance shall be provided on the situations in which the use of

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

					the feed is appropriate. When the feed corresponds to a significant part of the daily ration, guidance should be provided to prevent the risk of abrupt changes in the nature of the feed. Indicate on the instructions for use: Water should be available at all times.
[^{F1} Support the preparation for and recovery from sport effort	High level of selenium and a minimum content of 50 mg vitamin E per kg complete feed with moisture content of 12 %. The complementary feed may contain compounds of selenium in a concentration higher than 100 times the relevant fixed maximum content in complete feed.	Equidae	Total amount of — vitamin E — selenium	Up to 8 weeks before sport effort – Up to 4 weeks after sport effort	The instructions for use for the feed shall ensure that the legal maximum contents of selenium for complete feed are respected.]

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

Nutritional restoration, convalescence	High concentration of essential nutrients and highly digestible ingredients	Equines	—	Highly digestible ingredients including their treatment if appropriate Content of n-3 and n-6 fatty acids (if added)	Until restoration is achieved	Guidance shall be provided on the situations in which the use of the feed is appropriate. In the case of feedingstuffs specially presented to be given via tubing, indicate on the package, container or label: Administration under veterinary supervision.
Support of liver function in the case of chronic liver insufficiency	Low level of protein but of high quality and highly digestible carbohydrates	Equines	— — — —	Protein and fibre sources Highly digestible carbohydrates including their treatment if appropriate Methionine Choline Content of n-3 fatty acids (if added)	Initially up to six months	Guidance should be provided on the manner in which the feed should be fed including many small meals per day. Indicate on the package, container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use.
Support of renal function in the case of	Low level of protein but of high quality	Equines	— —	Protein sources Calcium	Initially up to (s) months	Indicate on the package,

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

chronic renal insufficiency	and low level of phosphorus	— — — —	Phosphorus Potassium Magnesium Sodium	container or label: It is recommended that a veterinarian's opinion be sought before use or before extending the period of use. Indicate in the instructions for use: Water should be available at all times.
a	[^{F2} If appropriate the manufacturer may also recommend use for temporary renal insufficiency.			
b	If the feed is recommended for temporary renal insufficiency the recommended period for use shall be two to four weeks.]			
c	In the case of feedingstuffs for cats, 'feline lower urinary tract disease' or 'feline urological syndrome — FUS' may complete the particular nutritional purpose.			
d	In the case of feedingstuffs for a particular intolerance reference to the specific intolerance can replace 'ingredient and nutrient'.			
e	The manufacturer may complete the particular nutritional purpose with the reference: 'exocrine pancreatic insufficiency'.			
f	[^{F3} The manufacturer may complete the particular nutritional purpose with a reference to 'Feline hepatic lipidosis'.]			
g	The term 'ketosis' may be replaced by 'acetonaemia'.			
h	The manufacturers may also recommend the use for ketosis recuperation.			
i	In the case of feedingstuffs for dairy cows.			
j	In the case of feedingstuffs for ewes.			
k	In the case of feedingstuffs for dairy cows: 'maximum two months from the start of lactation'.			
l	Indicate the category of ruminants concerned.			
m	[^{F4} Calculated by the Strong Ion Difference method (SID-Value): SID is the difference between the sums of concentrations of the strong cations and strong anions; [SID] = [mmol Na ⁺ /l] + [mmol K ⁺ /l] + [mmol Ca ⁺⁺ /l] + [mmol Mg ⁺⁺ /l] – [mmol Cl/l] – [mmol Other Strong Anions/l].]			
n	In the case of feedingstuffs specially prepared to meet the specific conditions of very old animals (easily ingestible ingredients) a reference to 'old animals' shall complete the indication of the species or category of animals.			

Textual Amendments

- F2** Substituted by [Commission Regulation \(EU\) No 1123/2014 of 22 October 2014 amending Directive 2008/38/EC establishing a list of intended uses of animal feedingstuffs for particular nutritional purposes \(Text with EEA relevance\)](#).
- F3** Substituted by [Commission Regulation \(EU\) No 5/2014 of 6 January 2014 amending Directive 2008/38/EC establishing a list of intended uses of animal feedingstuffs for particular nutritional purposes \(Text with EEA relevance\)](#).

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

- F4** Inserted by Commission Regulation (EU) No 1123/2014 of 22 October 2014 amending Directive 2008/38/EC establishing a list of intended uses of animal feedingstuffs for particular nutritional purposes (Text with EEA relevance).
- F5** Inserted by Commission Regulation (EU) No 1070/2010 of 22 November 2010 amending Directive 2008/38/EC by adding to the list of intended uses as a particular nutritional purpose the support of the metabolism of the joints in the case of osteoarthritis in dogs and cats (Text with EEA relevance).

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

- (1) [OJ L 268, 18.10.2003, p. 29.](#)
- (2) [OJ L 270, 14.12.1970, p. 1.](#)