

## ANNEX VI

Reference methods for assessment of concentrations of sulphur dioxide, nitrogen dioxide and oxides of nitrogen, particulate matter (PM<sub>10</sub> and PM<sub>2,5</sub>), lead, benzene, carbon monoxide, and ozone

[<sup>F1</sup>A. Reference methods for the assessment of concentrations of sulphur dioxide, nitrogen dioxide and oxides of nitrogen, particulate matter (PM<sub>10</sub> and PM<sub>2,5</sub>), lead, benzene, carbon monoxide and ozone

1. Reference method for the measurement of sulphur dioxide

The reference method for the measurement of sulphur dioxide is that described in EN 14212:2012 ‘Ambient air — Standard method for the measurement of the concentration of sulphur dioxide by ultraviolet fluorescence’.

2. Reference method for the measurement of nitrogen dioxide and oxides of nitrogen

The reference method for the measurement of nitrogen dioxide and oxides of nitrogen is that described in EN 14211:2012 ‘Ambient air — Standard method for the measurement of the concentration of nitrogen dioxide and nitrogen monoxide by chemiluminescence’.

[<sup>X1</sup>3. Reference method for the sampling and measurement of lead

The reference method for the sampling of lead is that described in Section A(4) of this Annex. The reference method for the measurement of lead is that described in EN 14902:2005 ‘Standard method for measurement of Pb/Cd/As/Ni in the PM<sub>10</sub> fraction of suspended particulate matter’.]

**Editorial Information**

- X1** Substituted by [Corrigendum to Commission Directive \(EU\) 2015/1480 of 28 August 2015 amending several annexes to Directives 2004/107/EC and 2008/50/EC of the European Parliament and of the Council laying down the rules concerning reference methods, data validation and location of sampling points for the assessment of ambient air quality \(Official Journal of the European Union L 226 of 29 August 2015\)](#).

4. Reference method for the sampling and measurement of PM<sub>10</sub>

The reference method for the sampling and measurement of PM<sub>10</sub> is that described in EN12341:2014 ‘Ambient Air — standard gravimetric measurement method for the determination of the PM<sub>10</sub> or PM<sub>2,5</sub> mass concentration of suspended particulate matter’.

5. Reference method for the sampling and measurement of PM<sub>2,5</sub>

The reference method for the sampling and measurement of PM<sub>2,5</sub> is that described in EN12341:2014 ‘Ambient Air — standard gravimetric measurement method for the determination of the PM<sub>10</sub> or PM<sub>2,5</sub> mass concentration of suspended particulate matter’

[<sup>X1</sup>6. Reference method for the sampling and measurement of benzene

The reference method for the measurement of benzene is that described in EN 14662:2005, parts 1, 2 and 3 ‘Ambient air quality — Standard method for measurement of benzene concentrations’.]

7. Reference method for the measurement of carbon monoxide

---

*Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.*

---

The reference method for the measurement of carbon monoxide is that described in EN 14626:2012 ‘Ambient air — Standard method for the measurement of the concentration of carbon monoxide by non-dispersive infrared spectroscopy’.

8. Reference method for measurement of ozone

The reference method for the measurement of ozone is that described in EN 14625:2012 ‘Ambient air — Standard method for the measurement of the concentration of ozone by ultraviolet photometry’.]