

**COMMISSION REGULATION (EC) No 1248/2001****of 22 June 2001****amending Annexes III, X and XI to Regulation (EC) No 999/2001 of the European Parliament and of the Council as regards epidemio-surveillance and testing of transmissible spongiform encephalopathies**

THE COMMISSION OF THE EUROPEAN COMMUNITIES,

Having regard to the Treaty establishing the European Community,

Having regard to Regulation (EC) No 999/2001 of the European Parliament and of the Council of 22 May 2001 laying down rules for the prevention, control and eradication of certain transmissible spongiform encephalopathies <sup>(1)</sup>, and in particular Articles 20(2) and 23 thereof,

Whereas:

- (1) Detailed rules for monitoring of transmissible spongiform encephalopathies (TSEs) in bovine, ovine and caprine animals are laid down in Annex III to Regulation (EC) No 999/2001. These rules include systematic testing of bovine animals over 30 months of age entering the food chain and random testing of bovine animals over 30 months of age not entering the food chain. In addition, all bovine animals subject to casualty slaughter or found sick when slaughtered for destruction under the over thirty months scheme (OTMS) shall be tested. Ovine and caprine animals with clinical signs compatible with TSE shall be subject to active surveillance.
- (2) In view of the detection of bovine spongiform encephalopathy (BSE) in two bovine animals at the age of 28 months in routine testing of casualty slaughtered animals and to provide an early warning system of the emergence of any unfavourable trends in BSE incidence in younger animals, the age limit should be reduced to 24 months in animals belonging to certain risk populations.
- (3) In the monitoring carried out during the first trimester of 2001, positive BSE cases were detected in all Member States except in Greece, Luxembourg, Austria, Finland and Sweden. The number of bovine animals belonging to certain risk groups tested in the above Member States were: 248 in Greece, 763 in Luxembourg, 3 295 in Austria, 4 527 in Finland and 8 254 in Sweden.
- (4) In its opinion of 6 July 2000 on the geographical risk of BSE (GBR), the Scientific Steering Committee (SSC) concluded that the GBR level of Luxembourg was III (BSE confirmed at a low level) and the GBR level of Austria, Finland and Sweden was II (BSE unlikely, but

not excluded). Greece did not submit a dossier for assessment, referring to legal and technical uncertainties.

- (5) In the light of the monitoring carried out in Austria, Finland and Sweden and of the assessment of the SSC, the presence of BSE in those Member States is unlikely, but not excluded. If present, BSE would most likely be detected by examining bovine animals which have died on farms, which have been subject to casualty slaughter or have been found sick at normal slaughter. Those Member States should therefore be allowed to reduce the testing in healthy slaughtered bovine animals.
- (6) With a view to additional information of the occurrence of BSE in the United Kingdom, the testing in the OTMS should be expanded to include all animals born within one year after the effective enforcement of the feed ban. Other bovine animals slaughtered under the OTMS should be tested at random.
- (7) Member States should be allowed to test other bovine animals on a voluntary basis, in particular where those animals are considered to present a higher risk, provided it is done without disrupting trade.
- (8) It is necessary to clarify the measures following testing of bovine animals and to introduce measures to prevent carcasses potentially contaminated by test-positive carcasses from entering the food chain.
- (9) Rapid post-mortem testing should be introduced on a random basis to improve the detection of scrapie in ovine and caprine animals. To obtain a more complete picture of the situation, it is necessary to carry out random sampling in two different target populations: dead-on-farm animals and slaughtered animals.
- (10) In Member States with small national herds of sheep and goats it is difficult to carry out statistically meaningful sampling in both target groups. Those Member States should therefore be allowed to use a smaller sample size but targeted at animals where the likelihood to find positive cases is highest.
- (11) In view of the role of genetic resistance in the development of clinical scrapie and the possibility to use breeding programmes in the prevention, control and eradication of scrapie, the genotype of all scrapie cases should be determined and cases found in resistant genotypes should be submitted for strain-typing.

<sup>(1)</sup> OJ L 147, 31.5.2001, p. 1.

- (12) The list of national reference laboratories should be updated.
- (13) Following the introduction of rapid tests in monitoring programmes for ovine and caprine animals it is necessary to lay down appropriate diagnostic methods and protocols. Furthermore, the diagnostic methods and protocols laid down for bovine animals should be updated.
- (14) According to Article 22 of Regulation (EC) No 999/2001, a conclusive statistical survey shall be used to confirm or overturn the risk analysis conclusions carried out as the first step in determining the BSE status of a country or a region. The minimum criteria for the statistical survey are laid down in Part B of Annex XI. In view of the lower BSE risk in Austria, Finland and Sweden, as assessed by the SSC, and the disproportionate resources involved, a derogation should be provided for those Member States to exclude from the survey dead-on-farm animals in remote areas with a low animal density.
- (15) In the interest of clarity Commission Decisions 98/272/EC<sup>(1)</sup> on epidemio-surveillance for transmissible spongiform encephalopathies, as last amended by Decision 2001/8/EC<sup>(2)</sup>, and 2000/764/EC<sup>(3)</sup> on the testing of bovine animals for the presence of bovine spongiform encephalopathy, as amended by Decision 2001/8/EC, should be repealed.
- (16) The measures provided for in this Regulation are in accordance with the opinion of the Standing Veterinary Committee,

HAS ADOPTED THIS REGULATION:

*Article 1*

Regulation (EC) No 999/2001 is amended as follows:

1. The text in Annex III is replaced by the text in Annex I to this Regulation.
2. The text in Annex X, Chapter A, point 3, is replaced by the text in Annex II to this Regulation.
3. The text in Annex X, Chapter C, is replaced by the text in Annex III to this Regulation.
4. The text in Annex XI, Chapter B, is replaced by the text in Annex IV to this Regulation.

*Article 2*

1. Decisions 98/272/EC and 2000/764/EC are repealed.
2. References to the repealed Decisions shall be construed as references to Regulation (EC) No 999/2001. In particular, references to Annex IVA of Decision 98/272/EC shall be construed as references to Annex X, Chapter C, point 4, to Regulation (EC) No 999/2001.

*Article 3*

This Regulation shall enter into force on the day following its publication in the *Official Journal of the European Communities*.

It shall apply from 1 July 2001. However, the provisions of Annex III, Chapter A, section II, to Regulation (EC) No 999/2001, as set out in Annex I to this Regulation, shall apply from 1 January 2002.

The provisions of Annex III to Regulation (EC) No 999/2001, as set out in Annex I to this Regulation, shall be reviewed in the light of the results obtained during the first six months of the monitoring.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 22 June 2001.

*For the Commission*

David BYRNE

*Member of the Commission*

<sup>(1)</sup> OJ L 122, 24.4.1998, p. 59.

<sup>(2)</sup> OJ L 2, 5.1.2001, p. 28.

<sup>(3)</sup> OJ L 305, 6.12.2000, p. 28.

## ANNEX I

## 'ANNEX III

## MONITORING SYSTEM

## CHAPTER A

## I. MONITORING IN BOVINE ANIMALS

## 1. General

Monitoring in bovine animals shall be carried out in accordance with the laboratory methods laid down in Annex X, Chapter C, point 3.1(b).

## 2. Monitoring in animals slaughtered for human consumption

## 2.1. All bovine animals over 24 months of age:

- subject to "special emergency slaughtering" as defined in Article 2(n) of Council Directive 64/433/EEC <sup>(1)</sup>, or
  - slaughtered in accordance with Annex I, Chapter VI, point 28(c), to Directive 64/433/EEC,
- shall be tested for BSE.

## 2.2. All bovine animals over 30 months of age subject to normal slaughter for human consumption shall be tested for BSE.

## 2.3. By way of derogation from point 2.2, and with regard to bovine animals born, reared and slaughtered on their territory, Austria, Finland and Sweden may decide to examine only a random sample. The sample shall comprise at least 10 000 animals per year.

## 3. Monitoring in animals not slaughtered for human consumption

Bovine animals over 24 months of age which have died or been killed but which were not

- killed for destruction pursuant to Commission Regulation (EC) No 716/962 <sup>(2)</sup>,
- killed in the framework of an epidemic, such as foot-and-mouth disease,
- slaughtered for human consumption,

shall be tested for BSE at random. The number of samples shall not be less than the sample size indicated in the table. The sampling must be representative for each region and continuous.

Total population over 24 months	Sample size (*)	Total population over 24 months	Sample size (*)
100 000	950	4 500 000	6 000
200 000	1 550	5 000 000	6 500
300 000	1 890	5 500 000	7 000
400 000	2 110	6 000 000	7 500
500 000	2 250	6 500 000	8 000
600 000	2 360	7 000 000	8 500
700 000	2 440	7 500 000	9 000
800 000	2 500	8 000 000	9 500
900 000	2 550	8 500 000	10 000
1 000 000	2 590	9 000 000	10 500
1 500 000	3 000	9 500 000	11 000
2 000 000	3 500	10 000 000	11 500
2 500 000	4 000	10 500 000	12 000
3 000 000	4 500	11 000 000	12 500
3 500 000	5 000	11 500 000	13 000
4 000 000	5 500	12 000 000	13 500

(\*) The sample size has been calculated to detect a prevalence of 0,1 % with a 95 % confidence in the subpopulation referred to in point 3, based on the assumption that the proportion of this subpopulation in the total population of bovine animals over 24 months of age is 1 %. Where the size of the total population of bovine animals over 24 months of age is 1 500 animals or more, the sample size has been increased by 500 samples per 500 000 animals as a proportionality adjustment, to take account of the larger likelihood of variation in risk for BSE within the population.

<sup>(1)</sup> OJ 121, 29.7.1964, p. 2012/64.

<sup>(2)</sup> OJ L 99, 20.4.1996, p. 14.

#### 4. **Monitoring in animals purchased for destruction pursuant to Regulation (EC) No 716/96**

- 4.1. All animals subject to casualty slaughter or found sick at ante-mortem inspection shall be tested for BSE.
- 4.2. All animals born between 1 August 1996 and 1 August 1997 shall be tested for BSE.
- 4.3. A random sample comprising at least 50 000 animals annually of animals not covered by points 4.1 or 4.2 shall be tested for BSE.

#### 5. **Monitoring in other animals**

In addition to the testing referred to in points 2 to 4, Member States may on a voluntary basis decide to test other bovine animals on their territory, in particular where those animals originate from countries with indigenous BSE, have consumed potentially contaminated feedingstuffs or were born or derived from BSE infected dams.

#### 6. **Measures following testing**

- 6.1. Where an animal slaughtered for human consumption is tested for BSE, the health marking provided for in Chapter XI of Annex I to Directive 64/433/EEC shall not be carried out on the carcase of that animal until a negative result to the rapid test has been obtained.
- 6.2. Member States may derogate from the provisions of point 6.1 where an official system is in place in the slaughterhouse ensuring that no parts of examined animals bearing the health mark leave the slaughterhouse until a negative result to the rapid test has been obtained.
- 6.3. All parts of the body of an animal tested for BSE including the hide shall be retained under official control until a negative result to the rapid test has been obtained, unless they are destroyed in accordance with Annex V, point 3 or 4.
- 6.4. All parts of the body of an animal found positive to the rapid test including the hide shall be destroyed in accordance with Annex V, point 3 or 4, apart from material to be retained in conjunction with the records provided for in Chapter B, Section III.
- 6.5. Where an animal slaughtered for human consumption is found positive to the rapid test, at least the carcase immediately preceding the test-positive carcase and two carcasses immediately following the test-positive carcase on the same slaughter line shall be destroyed in accordance with point 6.4, in addition to the test-positive carcase.
- 6.6. Member States may derogate from the provisions of point 6.5 where a system is in place in the slaughterhouse preventing contamination between carcasses.

## II. MONITORING IN OVINE AND CAPRINE ANIMALS

### 1. **General**

Monitoring in ovine and caprine animals shall be carried out in accordance with the laboratory methods laid down in Annex X, Chapter C, point 3.2(b).

### 2. **Monitoring in animals slaughtered for human consumption**

Animals over 18 months of age slaughtered for human consumption shall be tested in accordance with the sample size indicated in the table. The sampling shall be representative for each region and season. The sample selection shall be designed with a view to avoid the over-representation of any group as regards the origin, species, age, breed, production type or any other characteristic. The age of the animals shall be estimated based on dentition, obvious signs of maturity or other reliable information. Multiple sampling in the same flock shall be avoided, where possible.

Total number of slaughtered animals over 18 months of age	Minimum sample size, slaughtered animals (*)	Total number of slaughtered animals over 18 months of age	Minimum sample size, slaughtered animals (*)
5 000	4 750	60 000	13 260
10 000	7 760	70 000	13 490
15 000	9 470	80 000	13 660
20 000	10 540	90 000	13 800
25 000	11 270	100 000	13 910
30 000	11 790	150 000	14 250
40 000	12 490	200 000	14 430
50 000	12 940		

Total number of slaughtered animals over 18 months of age	Minimum sample size, slaughtered animals (*)	Total number of slaughtered animals over 18 months of age	Minimum sample size, slaughtered animals (*)
250 000	14 540	1 100 000	14 880
300 000	14 610	1 200 000	14 890
350 000	14 660	1 300 000	14 890
400 000	14 700	1 400 000	14 900
450 000	14 730	1 500 000	14 900
500 000	14 760	1 600 000	14 910
600 000	14 790	1 700 000	14 910
700 000	14 820	1 800 000	14 920
800 000	14 840	1 900 000	14 920
900 000	14 850	2 000 000	14 920
1 000 000	14 870	2 100 000	14 920
		2 200 000 or more	14 930

(\*) The sample size has been calculated to detect a prevalence of 0,02 % with a 95 % confidence in slaughtered animals.

### 3. Monitoring in animals not slaughtered for human consumption

Animals over 18 months of age which have died or been killed, but which were not:

- killed in the framework of an epidemic, such as foot-and-mouth disease,
- slaughtered for human consumption,

shall be tested in accordance with the sample size indicated in the table. The sampling shall be representative for each region and season. The sample selection shall be designed with a view to avoid the over-representation of any group as regards the origin, species, age, breed, production type or any other characteristic. The age of the animal shall be estimated based on dentition, obvious signs of maturity or other reliable information. Multiple sampling in the same flock shall be avoided, where possible.

Total number of animals over 18 months of age (*)	Minimum sample size, dead animals (**)
100 000	950
200 000	1 550
300 000	1 890
400 000	2 110
500 000	2 250
600 000	2 360
700 000	2 440
800 000	2 500
900 000	2 550
1 000 000	2 590
1 500 000 or more	3 000

(\*) Where the total number of ovine and caprine animals over 18 months of age is not known, the total number of "ewes and ewe-lambs put to the ram" and "goats which have already kidded and goats mated" shall be used instead.

(\*\*) The sample size has been calculated to detect a prevalence of 0,1 % with a 95 % confidence in dead animals, based on the assumption that the proportion of dead animals in the total population of ovine and caprine animals over 18 months of age is 1 %.

#### 4. Monitoring in Member States with a small ovine and caprine population

Member States, where the total number of ovine and caprine animals over 18 months of age is 500 000 or less may, by way of derogation from the sampling foreseen in points 2 and 3, decide to monitor the following combined subpopulation:

- (a) animals over 18 months of age which have died or been killed, but which were not:
  - killed in the framework of an epidemic, such as foot-and-mouth disease,
  - slaughtered for human consumption ("dead animals"); and
- (b) animals over 18 months of age whose appearance suggests a chronic wasting condition ("chronic wasting animals").

The number of samples tested annually in each Member State from the above combined subpopulation shall not be less than the sample size indicated in the table.

The sampling shall be representative for each region and season. The sample selection shall be designed with a view to avoid the over-representation of any group as regards the origin, species, age, breed, production type or any other characteristic. Where dead animals are sampled, the age of the animal shall be estimated based on dentition, obvious signs of maturity or other reliable information. Where chronic wasting animals are sampled, only animals inspected by an official veterinarian and for which the age and clinical symptoms are well documented shall be selected for sampling. Multiple sampling in the same flock shall be avoided, where possible.

Total number of ovine and caprine animals over 18 months of age (*)	Minimum sample size, dead and chronic wasting animals
10 000	100
20 000	200
30 000	300
40 000	400
50 000	500
60 000	600
70 000	700
80 000	800
90 000	900
100 000	950
200 000	1 550
300 000	1 890
400 000	2 110
500 000	2 250

(\*) Where the total number of ovine and caprine animals over 18 months of age is not known, the total number of "ewes and ewe-lambs put to the ram" and "goats which have already kidded and goats mated" shall be used instead.

#### 5. Monitoring in other animals

In addition to the monitoring programmes set out in points 2 to 4, Member States may on a voluntary basis carry out monitoring in other animals, in particular:

- animals used for dairy production,
- animals originating from countries with indigenous TSEs,
- animals which have consumed potentially contaminated feedingstuffs,
- animals born or derived from TSE infected dams,
- animals from flocks infected with TSE.

#### 6. Measures following testing of ovine and caprine animals

All parts of the body of a tested animal including the hide shall be retained under official control until a negative result to the rapid test has been obtained, unless they are destroyed in accordance with Annex V, point 3 or 4.

All parts of the body of an animal found positive to the rapid test including the hide shall be destroyed in accordance with Annex V, point 3 or 4, apart from material to be retained in conjunction with the records provided for in Chapter B, section III.

## 7. Genotyping

The prion protein genotype shall be determined for each positive TSE case in sheep. TSE cases found in resistant genotypes (sheep of genotypes which encode alanin on both alleles at codon 136, arginin on both alleles at codon 154 and arginin on both alleles at codon 171) shall immediately be reported to the Commission. Where possible, such cases shall be submitted for strain-typing. Where strain-typing of such cases is not possible, the herd of origin and all other herds where the animal has been shall be subjected to enhanced monitoring with a view to find other TSE cases for strain-typing.

## CHAPTER B

### I. INFORMATION TO BE PRESENTED BY MEMBER STATES IN THEIR REPORT

1. The number of suspected cases per animal species placed under movement restrictions in accordance with Article 12(1).
2. The number of suspected cases per animal species subject to laboratory examination in accordance with Article 12(2) and the outcome of the examination.
3. The number of flocks where suspected cases in ovine and caprine animals have been reported and investigated pursuant to Article 12(1) and (2).
4. The estimated size of each subpopulation referred to in Chapter A, Section I, points 3 and 4.
5. The number of bovine animals tested within each subpopulation referred to in Chapter A, Section I, point 2 to 5, the method for sample selection and the outcome of the tests.
6. The estimated size of those subpopulations referred to in Chapter A, Section II, points 2 to 4, which have been selected for sampling.
7. The number of ovine and caprine animals and flocks tested within each subpopulation referred to in Chapter A, Section II, points 2 to 5, the method for sample selection and the outcome of the tests.
8. Number, age distribution and geographical distribution of positive cases of BSE and scrapie. The country of origin, if not the same as the reporting country, of positive cases of BSE and scrapie. Number and geographical distribution of scrapie positive flocks. The year and, where possible, month of birth should be given for each BSE case.
9. Positive TSE cases confirmed in animals other than bovine, ovine and caprine animals.

### II. INFORMATION TO BE PRESENTED BY THE COMMISSION IN ITS SUMMARY

The summary shall be presented in a tabled format covering at least the information referred to in Part I for each Member State.

### III. RECORDS

1. The competent authority shall keep, for seven years, records of:
  - the number and types of animals placed under movement restrictions as referred to in Article 12(1),
  - the number and outcome of clinical and epidemiological investigations as referred to in Article 12(1),
  - the number and outcome of laboratory examinations as referred to in Article 12(2),
  - the number, identity and origin of animals sampled in the framework of the monitoring programmes as referred to in Chapter A and, where possible, age, breed and anamnestic information,
  - the prion protein genotype of positive TSE cases in sheep,
  - where chronic wasting ovine and caprine animals have been selected for sampling, the method to establish the age of and the clinical symptoms observed in each sampled animal.
2. The investigating laboratory shall keep, for seven years, all records of testing, in particular laboratory workbooks and, where appropriate, paraffin blocks and photographs of Western blots.

## ANNEX II

3. The national reference laboratories are:

Austria:	Bundesanstalt für Tierseuchenbekämpfung, Mödling Robert Koch Gasse 17 A-2340 Mödling	
Belgium:	CERVA-CODA-VAR Centre d'Étude et de Recherches Vétérinaires et Agrochimiques Centrum voor Onderzoek in Diergeneeskunde en Agrochemie Veterinary and Agrochemical Research Centre Groeselenberg 99 B-1180 Bruxelles	
Denmark:	Danish Veterinary Laboratory Bülowsvej 27 DK-1790 Copenhagen V	
Finland:	Eläinlääkintä- ja elintarvikelaitos Hämeentie 57 FIN-00550 Helsinki	
France:	Agence Française de Sécurité Sanitaire des Aliments Laboratoire de pathologie bovine 31, avenue Tony Garnier BP 7033 F-69342 Lyon Cedex	
Germany:	Bundesforschungsanstalt für Viruskrankheiten der Tiere Anstaltsteil Insel Riems Boddenblick 5A D-17498 Insel Riems	
Greece:	Laboratory of Microbiology and Infectious Diseases Faculty of Veterinary Medicine Aristotelian University of Thessaloniki University Campus GR-54006 Thessaloniki (rapid and immunological tests)	Laboratory of Gross Pathology (Morgue) Faculty of Veterinary Medicine Aristotelian University of Thessaloniki Giannitson & Voutyra St GR-54627 Thessaloniki (histopathology)
Ireland:	Central Veterinary Research Laboratory Abbotstown Castleknock Dublin 15 Ireland	
Italy:	Istituto Zooprofilattico Sperimentale del Piemonte Liguria e Valle d'Aosta CEA Via Bologna I-148-10150 Torino	
Luxembourg:	CERVA-CODA-VAR Centre d'Étude et de Recherches Vétérinaires et Agrochimiques Centrum voor Onderzoek in Diergeneeskunde en Agrochemie Veterinary and Agrochemical Research Centre Groeselenberg 99 B-1180 Bruxelles	
Netherlands:	Instituut voor Dierhouderij en Diergezondheid, ID-DLO Lelystad Edelhertweg 15 Postbus 65 8200 AB Lelystad Netherlands	



---

Portugal:	Laboratório Nacional de Investigação Veterinária Estrada de Benfica, 701 P-1500 Lisboa
Spain:	Laboratorio de la Facultad de Veterinaria Departamento de Patología Animal (Anatomía Patológica) Zaragoza Spain (BSE and scrapie, methods other than rapid tests)  Laboratorio Central de Veterinaria de Algete Madrid Spain (rapid tests)  Centro de Investigacion en Sanidad Animal (CISA) Ctra, De Algete al Casar de Talamanca 28130 Valdeolmos (Madrid) Spain (TSEs other than BSE or scrapie)
Sweden:	National Veterinary Institute S-751 89 Uppsala
United Kingdom	Veterinary Laboratories Agency Woodham Lane New Haw Addlestone Surrey KT15 3NB United Kingdom'

---

## ANNEX III

## CHAPTER C

**Sampling and laboratory testing****1. Sampling**

Any samples intended to be examined for the presence of a TSE shall be collected using the methods and protocols laid down in the latest edition of the *Manual of standards for diagnostic tests and vaccines* of the International Office for Epizootics (IOE/OIE) (hereinafter referred to as "the Manual"). In the absence of such methods and protocols, the samples shall be collected in a manner appropriate for the correct application of tests. The samples shall be correctly marked as to the identity of the sampled animal.

**2. Laboratories**

Any laboratory examination for TSE shall be carried out in laboratories approved for that purpose.

**3. Methods and protocols****3.1. Laboratory testing for the presence of BSE in bovine animals****(a) Suspect cases**

Tissues from bovine animals sent for laboratory testing pursuant to the provisions of Article 12(2) shall be subject to a histopathological examination as laid down in the latest edition of the Manual, except where the material is autolysed. Where the result of the histopathological examination is inconclusive or negative or where the material is autolysed, the tissues shall be subjected to an examination by one of the other diagnostic methods laid down in the Manual (immunocytochemistry, immuno-blotting or demonstration of characteristic fibrils by electron microscopy). However, rapid tests cannot be used for this purpose.

If the result of one of the above examinations is positive, the animals shall be regarded a positive BSE case.

**(b) BSE monitoring**

Tissues from bovine animals sent for laboratory testing pursuant to the provisions of Annex III, Chapter A, Section I (Monitoring in bovine animals) shall be examined by a rapid test.

When the result of the rapid test is inconclusive or positive, the tissues shall immediately be subject to confirmatory examinations in an official laboratory. The confirmatory examination shall start by a histopathological examination of the brainstem as laid down in the latest edition of the Manual, except where the material is autolysed or otherwise not suitable for examination by histopathology. Where the result of the histopathological examination is inconclusive or negative or where the material is autolysed, the tissues shall be subjected to an examination by one of the other diagnostic methods mentioned under (a).

An animal shall be regarded a positive BSE case if the result of the rapid test is positive or inconclusive, and

- the result of the subsequent histopathological examination is positive, or
- the result of another diagnostic method mentioned under (a) is positive.

**3.2. Laboratory testing for the presence of scrapie in ovine and caprine animals****(a) Suspect cases**

Tissues from ovine and caprine animals sent for laboratory testing pursuant to the provisions of Article 12(2) shall be subject to a histopathological examination as laid down in the latest edition of the Manual, except where the material is autolysed. Where the result of the histopathological examination is inconclusive or negative or where the material is autolysed, the tissues shall be subjected to an examination by immunocytochemistry or immuno-blotting, as laid down in the Manual. However, rapid tests cannot be used for this purpose.

If the result of one of the above examinations is positive, the animal shall be regarded a positive scrapie case.

**(b) Scrapie monitoring**

Tissues from ovine and caprine animals sent for laboratory testing pursuant to the provisions of Annex III, Chapter A, Section II (Monitoring in ovine and caprine animals) shall be examined by a rapid test.

When the result of the rapid test is inconclusive or positive, the brainstem shall immediately be sent to an official laboratory for confirmatory examinations by immunocytochemistry or immuno-blotting, as referred to under (a).

An animal shall be regarded a positive scrapie case if the result of the confirmatory examination is positive.

3.3. *Laboratory testing for the presence of TSEs other than those referred to in points 3.1 and 3.2*

The tests carried out to confirm the suspected presence of a TSE different from those referred to in points 3.1 and 3.2 shall include at least a histopathological examination of brain tissue. The competent authority may also require laboratory tests such as immunocytochemistry, immuno-blotting, demonstration of characteristic fibrils by electron microscopy or other methods designed to detect the disease associated form of the prion protein. In any case at least one other laboratory examination shall be carried out if the initial histopathological examination is negative or inconclusive. At least three different examinations shall be carried out in the event of the first appearance of the disease.

In particular, where BSE is suspected in a species other than bovine animals, samples shall be submitted for strain-typing, where possible.

4. **Rapid tests**

For the purposes of carrying out the tests in accordance with Article 5(3) and Article 6(1), the following methods shall be used as rapid tests within the meaning of this Regulation:

- immuno-blotting test based on a western blotting procedure for the detection of the protease-resistant fragment PrP<sup>Res</sup> (Prionics Check test),
- chemiluminescent ELISA test involving an extraction procedure and an ELISA technique, using an enhanced chemiluminescent reagent (Enfer test),
- sandwich immunoassay for PrP<sup>Res</sup> carried out following denaturation and concentration steps (Bio-Rad Platelia test).

5. **Alternative tests**

(To be defined)

---

ANNEX IV

**B. Concerning statistical surveys**

1. The statistical survey referred to in Article 22 must cover:
  - the animals sampled pursuant to the provisions of Annex III, Chapter A, Section I, points 2.1 and 4.1,
  - all the animals in the subpopulation referred to in Annex III, Chapter A, Section I, point 3, instead of a random sample.

This provision, applicable for a period of one year, may be reviewed in the light of experience gained in the first six months.

2. Austria, Finland and Sweden may decide to derogate from the provisions of point 1, second indent, in remote areas with a low animal density.'
-