

▼B

COMMISSION REGULATION (EC) No 1293/2008

of 18 December 2008

concerning the authorisation of a new use of *Saccharomyces cerevisiae* CNCM I-1077 (Levucell SC20 and Levucell SC10 ME) as a feed additive

(Text with EEA relevance)

Article 1

The preparation specified in the Annex, belonging to the additive category 'zootechnical additives' and to the functional group 'gut flora stabilisers', is authorised as an additive in animal nutrition subject to the conditions laid down in that Annex.

Article 2

This Regulation shall enter into force on the 20th day following its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

ANNEX

Identification number of the additive	Name of the holder of authorisation	Additive (trade name)	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
						CFU/kg of complete feedingstuff with a moisture content of 12 %			
Category of zootechnical additives. Functional group: gut flora stabilisers									
4b1711	►M2 Danstar Ferment AG represented by Lallemand SAS ◀	<i>Saccharomyces cerevisiae</i> CNCM I-1077 (Levucell SC20, Levucell SC10 ME)	<p>Additive composition:</p> <p>Solid form:</p> <p>Preparation of <i>Saccharomyces cerevisiae</i> CNCM I-1077 of viable dried cells with a guaranteed minimal concentration of 2×10^{10} CFU/g.</p> <p>Coated form:</p> <p>Preparation of <i>Saccharomyces cerevisiae</i> CNCM I-1077 of viable dried cells with a guaranteed minimal concentration of 1×10^{10} CFU/g</p> <p>Characterisation of the active substance:</p> <p><i>Saccharomyces cerevisiae</i> CNCM I-1077: 80 % of viable dried cells and 14 % of not viable cells.</p> <p>Analytical method (1):</p> <p>Pour plate method and molecular identification (PCR).</p>	Lambs	—	$3,0 \times 10^9$	►M1 — ◀	<ol style="list-style-type: none"> 1. In the directions for use of the additive and premixtures, indicate the storage temperature, storage life, and stability to pelleting. 2. In complementary feeding-stuffs, do not exceed 50 °C with Levucell SC20 and 80 °C with Levucell SC10ME. 3. Coated form, only for inclusion through a pelleted feed. 4. Recommended dose: $7,3 \times 10^9$ CFU/kg of complete feedingstuff. 5. If the product is handled or mixed in confined atmosphere, it is recommended to use safety glasses and masks for mixing if the mixers are not equipped with exhaust systems. 	8.1.2019

(1) Details of the analytical methods are available at the following address of the Community Reference Laboratory: www.irmm.jrc.be/crl-feed-additives